

# POTATO

**AbdA™** - Improves vegetative growth and tuber development.

**AbdA™** - Increases the number of roots / tubers.

**AbdA™** - For uniform maturity and tuber size.

**All the above actions shall lead to a plentiful harvest**

## Dosage and Methods of Application

### Time of Application

Between  
0 - 15 days  
of planting

### Dosage

4 kg per acre

### Method of Application

Apply uniformly along planting rows by mixing with 6 - 8 kg of dry coarse sand / soil.

Irrigate the field after **AbdA™** application.

Parrys  
BIO

# AbdA™

Crop Vitaliser and Yield Enhancer

Granule



**E.I.D.PARRY(INDIA) LIMITED**

Bio Products Division  
'Dare House'

234, N.S.C. Bose Road, Chennai - 600 001, India  
Phone : 91-44-25306789 Fax : 91-44-25341609

[www.parrysaza.com](http://www.parrysaza.com)



MURUGAPPA  
GROUP

# AbdA™

Crop Vitaliser and Yield Enhancer

**Granule**

- AbdA™** - **AbdA™** is a resultant product of a decade of research and field trials of the Bio Products Division of E.I.D. Parry (India) Ltd.
- AbdA™** - Granulated herbal compound prepared by combining specific herbal extracts in proportionate ratio.
- AbdA™** - Application is easier due to granular form.
- AbdA™** - Environmental friendly, free of residual and side effects.

## Notable Beneficial Effects

- AbdA™** - Produces balanced root and shoot ratio for the initial establishment of the crop.
- AbdA™** - Gives an uniform and robust vegetative growth.
- AbdA™** - It's slow release mechanism keeps the crop healthy and greenish during the entire vegetative phase.
- AbdA™** - Induces insect pest and disease resistance.
- AbdA™** - Ensures uniform crop maturity.

**All the above actions shall lead to a plentiful harvest**

## RICE

- AbdA™** - Increases number of productive tillers.
- AbdA™** - The physiological activity in the crop continues till maturity.
- AbdA™** - Increases the luster and weight of grains.
- AbdA™** - Reduces the number of chaffy grains.

**All the above actions shall lead to a plentiful harvest**



Treated Untreated

Treated Untreated

## Dosage and Methods of Application

Crop Stage	Time of Application	Dosage	Method of Application
Nursery	Between 0 - 15 days of sowing	4 kg per acre	Apply by mixing with 6 - 8 kg dry coarse sand.
Main field	Between 0 - 15 days of transplanting	4 kg per acre	Apply by mixing with 6 - 8 kg dry coarse sand in transplanted field / puddle field.

**Caution :** There should be sufficient water in the field after **AbdA™** Application.

